

Quantitative approach to facilitated diffusion with intersegmental jumping

Michael A. Lomholt,¹ Bram van den Broek,^{2,3} Svenja-Marei J. Kalisch,² Gijs J.L. Wuite,² and Ralf Metzler⁴

¹*MEMPHYS - Center for Biomembrane Physics, Department of Physics and Chemistry, University of Southern Denmark, Campusvej 55, 5230 Odense M, Denmark*

²*Department of Physics and Astronomy, Faculty of Sciences, VU University, De Boelelaan 1081, 1081 HV Amsterdam, The Netherlands*

³*Physics of Life Processes, Leiden Institute of Physics, Leiden University, Niels Bohrweg 2, 2333 CA Leiden, The Netherlands*

⁴*Physics Department, Technical University of Munich, D-85747 Garching, Germany*

(Dated: December 16, 2008)

We study the impact of DNA coiling on the search rate of proteins moving along the DNA contour interspersed by three-dimensional (3D) bulk excursions. When the DNA is coiled proteins performing short 3D hops along a DNA segment can be captured by foreign DNA segments that have looped back close to the original segment. These intersegmental jumps are shown to enhance the search rate for a specific site on the DNA by lowering the tendency to resample previously visited sites on the DNA. The model developed here offers a quantitative description of recent single molecule experiments on facilitated diffusion of restriction enzymes EcoRV.

PACS numbers: 82.39.-k, 02.50.Ey

A single stochastic search mechanism is usually not efficient at optimizing a search process. Thus Brownian motion, whose fractal dimension is $d_f = 2$, in 1D and 2D leads to significant oversampling in the sense that a given site is revisited many times [1, 2]. Lévy motion with a jump length distribution $\lambda(x) \simeq |x|^{-1-\alpha}$ ($0 < \alpha < 2$) has $d_f = \alpha$ and thus a comparatively reduced degree of oversampling. However, the first arrival of Lévy motion is hampered by leapovers, i.e., frequent overshoots of the target [3]. In 3D both Brownian and Lévy motions are inefficient at locating a small target. While typical chemical reactants in a liquid environment have no choice but to passively diffuse until mutual encounter [4] intermittent models combine different search mechanisms [2, 5].

An important example for intermittent search is facilitated diffusion of DNA binding proteins searching for their specific target binding site on a DNA molecule. In the Berg-von Hippel model [6] proteins intermittently diffuse in 3D bulk or along the 1D DNA chain [7], the latter mediated by non-specific binding (NSB) [8].

Previous approaches to facilitated protein diffusion consider bulk exchange for a straight cylindrical chain [9, 10]; they consider scaling properties in different chain configurations [11] without quantitative comparison of the search rates between these; or they reformulate the problem with uncorrelated bulk excursions [12, 13, 14, 15]. Here we relate a multiparticle picture to single protein jumps along the DNA contour allowing us to consider DNA as a fluctuating chain. Comparison to recent single DNA measurements [16] demonstrates that our model quantitatively describes the effect of DNA coiling.

General model. In our description of the target search process, we use the density per length $n(x, t)$ of proteins on the DNA as the relevant dynamical quantity (x is the

distance along the DNA contour). We include 1D diffusion along the DNA with diffusivity D_{1d} , protein dissociation with rate $k_{\text{off}}^{\text{ns}}$ and (re)adsorption after diffusion through the bulk fluid with diffusion constant D_{3d} . The dynamics of $n(x, t)$ is thus governed by

$$\frac{\partial n(x, t)}{\partial t} = \left(D_{1d} \frac{\partial^2}{\partial x^2} - k_{\text{off}}^{\text{ns}} \right) n(x, t) - j(t) \delta(x) + G(x, t) + k_{\text{off}}^{\text{ns}} \int_{-\infty}^{\infty} dx' \int_0^t dt' W_{\text{bulk}}(x - x', t - t') n(x', t'). \quad (1)$$

Here, $j(t)$ is the flux into the target site located at $x = 0$, $W_{\text{bulk}}(x - x', t - t')$ is the joint probability that a protein returns to the point x at time t after leaving the DNA at x' , t' for a bulk excursion. $G(x, t)$ is the flux onto the DNA of proteins that have not previously been bound to the DNA. $j(t)$ is determined by imposing the boundary condition $n(x = 0, t) = 0$ at the target. To consider W_{bulk} as homogeneous in space and time we made two assumptions that we found to be valid for the system studied in [16]. First, that the DNA can be treated as being infinitely long when considering the search rate [17]. Second, that the coiled DNA conformation fluctuates quickly such that subsequent excursion distances $x - x'$ can be treated as independent.

To proceed, we Laplace and Fourier transform Eq. (1):

$$n(q, u)/W(q, u) - n_0(q) = G(q, u) - j(u) \quad (2)$$

with $n(q, u) = \mathcal{L}\{n(q, t)\}$; $n(q, t) = \mathcal{F}\{n(x, t)\}$; $n_0(x) = n(x, t = 0)$, and

$$W(q, u) = 1/\{u + D_{1d}q^2 + k_{\text{off}}^{\text{ns}}[1 - W_{\text{bulk}}(q, u)]\}. \quad (3)$$

We rewrite these equations in the form

$$n(q, u) = n^{\text{ns}}(q, u) - j(u)W(q, u) \quad (4)$$

where the density $n^{\text{ns}}(q, u) = W(q, u)[G(q, u) + n_0(q)]$ is the solution of the diffusion problem in absence of the target, i.e., when only NSB between proteins and the DNA occurs. In position space at $x = 0$ one obtains $j(u) = n^{\text{ns}}(x = 0, u)/W(x = 0, u)$.

In this study we focus on the long time behavior of $j(t)$ for which the density $n^{\text{ns}}(x, t)$ will reach an equilibrium value $n_{\text{eq}}^{\text{ns}}$. Since, by Tauberian theorems, large t correspond to small u we study the $u \rightarrow 0$ behavior of $j(u)$, with $n^{\text{ns}}(x, u) \sim n_{\text{eq}}^{\text{ns}}/u$. For $W_0(x = 0, u)$ we assume that the limit $W(x = 0, u = 0)$ is finite and non-zero; this is true when the distance distribution of bulk excursions is sufficiently long tailed as fulfilled for the W_{bulk} encountered below. In practice, being interested in the *first arrival* at the target, we assume that the protein concentration is sufficiently dilute such that this arrival happens after reaching $n_{\text{eq}}^{\text{ns}}$ ('rapid equilibrium') but still with many searching proteins. Thus $j(u) \sim k_{1d} n_{\text{eq}}^{\text{ns}}/u$, and therefore at long times the stationary current $j_{\text{stat}} \sim k_{1d} n_{\text{eq}}^{\text{ns}}$ into the target is reached. Here we introduced $k_{1d}^{-1} = W(x = 0, u = 0)$:

$$k_{1d}^{-1} = \int_{-\infty}^{\infty} \frac{dq}{2\pi} \frac{1}{D_{1d}q^2 + k_{\text{off}}(1 - \lambda_{\text{bulk}}(q))}. \quad (5)$$

$\lambda_{\text{bulk}}(x) = W_{\text{bulk}}(x, u = 0)$ is the distribution of relocation lengths along the DNA after a single 3D excursion.

We express our results in terms of the association rate

$$k_{\text{on}} = j_{\text{stat}}/n_{\text{bulk}} = k_{1d}K_{\text{ns}} \quad (6)$$

to the target where n_{bulk} is the density of unbound proteins in the bulk (at non-specific equilibrium). The constant of NSB per length of DNA is $K_{\text{ns}} = n_{\text{eq}}^{\text{ns}}/n_{\text{bulk}}$. In terms of the total volume concentration of proteins $n_{\text{total}} = n_{\text{bulk}} + l_{\text{DNA}}^{\text{total}} n_{\text{eq}}^{\text{ns}}$, where $l_{\text{DNA}}^{\text{total}}$ is the total length of DNA divided by the volume of the entire system, we have $j_{\text{stat}} \sim k_{\text{on}} n_{\text{total}}/(1 + K_{\text{ns}} l_{\text{DNA}}^{\text{total}})$. The rate k_{on} is related to the mean first arrival time for the steady state association. Namely $k_{\text{on}} n_{\text{bulk}}$ is the probability per time for protein association with the target. Thus the (survival) probability that no protein has arrived at the target yet is $P_{\text{surv}}(t) = \exp(-k_{\text{on}} n_{\text{bulk}} t)$, and the average target search time is $T = 1/(k_{\text{on}} n_{\text{bulk}})$. The flux $j(t)$ being linear in the protein concentration n_{total} implies that for sufficiently low n_{total} the steady state is reached well before the first protein actually binds to the specific target (note that *in vivo* protein concentrations can be as low as nanomolar).

Straight rod configuration (Fig. 1). Consider first stretched DNA corresponding to a cylinder of radius r_{int} (the range of non-specific interaction), with NSB reaction rate $k_{\text{on}}^{\text{ns}}$ at the boundary: the flux of proteins, per length of DNA, onto the DNA is $k_{\text{on}}^{\text{ns}}$ times the bulk concentration next to the DNA. This implies that at equilibrium of NSB, $n_{\text{eq}}^{\text{ns}} k_{\text{off}}^{\text{ns}} = n_{\text{bulk}} k_{\text{on}}^{\text{ns}}$ such that $K_{\text{ns}} = k_{\text{on}}^{\text{ns}}/k_{\text{off}}^{\text{ns}}$ [18].




FIG. 1: Schematic of the search mechanisms in Eq. (1) on a piece of stretched DNA. Note the tendency to revisit sites when immediate rebindings and thus short hops are likely.

To find the distribution $W_{\text{bulk}}(x, t)$ for return to the DNA of a protein released at the DNA at $x = 0$ and $t = 0$, we solve the cylindrical diffusion equation for the probability density $P(x, r, t)$ of the protein's position

$$\frac{\partial P}{\partial t} = D_{3d} \left(\frac{\partial^2}{\partial x^2} + \frac{1}{r} \frac{\partial}{\partial r} r \frac{\partial}{\partial r} \right) P \quad (7)$$

where r measures the distance to the cylinder axis. The reactive boundary condition for the probability flux out of the cylinder boundary is

$$-2\pi r_{\text{int}} D_{3d} \frac{\partial P}{\partial r} \Big|_{r=r_{\text{int}}} = -k_{\text{on}}^{\text{ns}} P|_{r=r_{\text{int}}} + \delta(t)\delta(x). \quad (8)$$

The initial distribution outside the cylinder is $P(r, t)|_{t=0} = 0$. From the solution of this problem we obtain W_{bulk} as $k_{\text{on}}^{\text{ns}} P|_{r=r_{\text{int}}}$ in terms of modified Bessel functions K_ν [with $\bar{q} = \sqrt{q^2 + u/D_{3d}}$]

$$W_{\text{bulk}}^{\text{cyl}}(q, u) = \left(1 + \frac{2\pi D_{3d} \bar{q} r_{\text{int}} K_1(\bar{q} r_{\text{int}})}{k_{\text{on}}^{\text{ns}} K_0(\bar{q} r_{\text{int}})} \right)^{-1}. \quad (9)$$

With these ingredients one can evaluate numerically the search rate to reach the specific target on a straight DNA, as given by Eq. (5). However, some limits allow analytic approximation for the integral in Eq. (5): for $k_{\text{on}}^{\text{ns}} \ll D_{3d}$ one may take $1 - \lambda_{\text{bulk}}(q) \approx 1$, and thus $k_{1d}^{-1} \sim (2\sqrt{D_{1d} k_{\text{off}}^{\text{ns}}})^{-1}$. One finds the association rate

$$k_{\text{on}} = 2l_{\text{sl}} k_{\text{on}}^{\text{ns}}, \quad (10)$$

where $l_{\text{sl}} = \sqrt{D_{1d}/k_{\text{off}}^{\text{ns}}}$ is the so-called sliding length of a protein during a single NSB period. Eq. (10) states that each time a protein, with rate constant $k_{\text{on}}^{\text{ns}}$, binds to the DNA within a distance l_{sl} from the target it is able to “slide” onto the target. Note that in this $k_{\text{on}}^{\text{ns}} \ll D_{3d}$ limit the search rate k_{on} is independent of the DNA conformation. In the opposite limit $k_{\text{on}}^{\text{ns}} \gg D_{3d}$ one may approximate $1 - \lambda_{\text{bulk}}(q) \approx 2\pi D_{3d}/[k_{\text{on}}^{\text{ns}}/\ln(l_{\text{sl}}^{\text{eff}}/r_{\text{int}})]$ where $l_{\text{sl}}^{\text{eff}} = \sqrt{k_{\text{on}}^{\text{ns}}/(2\pi D_{3d})} l_{\text{sl}}$. This produces the result [19]

$$k_{\text{on}} \sim 4\pi D_{3d} l_{\text{sl}}^{\text{eff}} / [\ln(l_{\text{sl}}^{\text{eff}}/r_{\text{int}})]^{1/2}. \quad (11)$$

In this limit the protein can rebind quickly after unbinding. This leads to oversampling and a lowered value of k_{on} compared to Eq. (10). $l_{\text{sl}}^{\text{eff}}$ can be interpreted as an effective sliding length including immediate rebindings




FIG. 2: Search rate enhancement on coiled DNA: a protein that in the case of straight DNA would perform a short hop can now, due to DNA looping, be captured by a foreign (remote) segment of DNA. This leads to a more efficient search since local oversampling is reduced by these intersegmental jumps [20]. The DNA configuration is assumed to be dynamic such that when the protein is released from the foreign segment the latter has moved further away from the original segment. Dashed lines: motion of the protein; dotted lines: DNA configurational changes.

[9]. Conversely, in the limit $k_{\text{on}}^{\text{ns}} \ll D_{3\text{d}}$ the rate of NSB will limit rebinding such that the protein will relocate by a long distance during a single 3D excursion. The part of λ_{bulk} corresponding to such long relocations do not contribute significantly to the integral in Eq. (5), which is the physical reason for the approximation leading to Eq. (10). Results (5) and (9) are identical to the classical result for straight, infinitely long DNA [9]. However, the approach here allows to explicitly consider coiled DNA.

Coil with persistence length (Fig. 2). To obtain W_{bulk} for a random coil DNA configuration we treat the DNA as consisting locally of straight segments. We assume that the local density l_{DNA} of DNA (length per volume) around a segment is dilute such that the typical distance $1/\sqrt{l_{\text{DNA}}}$ between segments is much larger than r_{int} . We then apply a superposition technique, where the probability that the protein has reacted with a segment is obtained by treating independently the interaction with each segment, and then choose among these the interaction for which binding happens first. To obtain the dynamics of the interaction of the protein with one segment of DNA we consider a large volume V around the segment of length L . Placing the protein at a uniformly random position in space relative to the DNA segment at $t = 0$ and denoting the probability that the protein has bound to this DNA before time t by $J_{\text{single}}(t)$, then the probability that it has not bound to any of N pieces of DNA is (using the assumed independent interactions) $P_{\text{surv}}^{\text{foreign}}(t) = [1 - J_{\text{single}}(t)]^N$. Taking the limit of $V, N, L \rightarrow \infty$ for fixed $l_{\text{DNA}} = NL/V$ we get $P_{\text{surv}}^{\text{foreign}}(t) = \exp[-J_{\text{cap}}(t)]$, where

$$\frac{u^2 J_{\text{cap}}(u)}{k_{\text{on}}^{\text{ns}} l_{\text{DNA}}} = \left(1 + \frac{k_{\text{on}}^{\text{ns}} K_0(\sqrt{u/D_{3\text{d}} r_{\text{int}}})}{2\pi \sqrt{u D_{3\text{d}} r_{\text{int}}} K_1(\sqrt{u/D_{3\text{d}} r_{\text{int}}})} \right)^{-1}. \quad (12)$$

W_{bulk} in the coiled conformation then becomes

$$W_{\text{bulk}}(q, t) = W_{\text{bulk}}^{\text{cyl}}(q, t) P_{\text{surv}}^{\text{foreign}}(t) - \frac{dP_{\text{surv}}^{\text{foreign}}(t)}{dt} \delta_{q,0} \int_t^\infty W_{\text{bulk}}^{\text{cyl}}(q=0, t') dt'. \quad (13)$$

The first term on the right hand side stands for proteins that return to the original segment without being captured by foreign segments. Those are represented by the second term, where the factor $\delta_{q,0}$ is a result of the assumption that a capture by foreign DNA segments leads to a long relocation measured along the DNA contour. These relocations only contribute close to $q = 0$ in Fourier space. Exploiting the reasoning behind the approximation that led to Eq. (10) we discard them from W_{bulk} when the integral in Eq. (5) is calculated. However, even numerically this integral is difficult to handle. A good approximation can be found noting that by Tauberian theorems the Laplace inversion for $J_{\text{cap}}(u)$ can be divided into three regimes when $k_{\text{on}}^{\text{ns}} \gg D_{3\text{d}}$:

$$J_{\text{cap}}(t) = \begin{cases} k_{\text{on}}^{\text{ns}} l_{\text{DNA}} t & , t \ll \frac{D_{3\text{d}} r_{\text{int}}^2}{(k_{\text{on}}^{\text{ns}})^2} \\ 4l_{\text{DNA}} r_{\text{int}} \sqrt{\pi D_{3\text{d}} t} & , \frac{D_{3\text{d}} r_{\text{int}}^2}{(k_{\text{on}}^{\text{ns}})^2} \ll t \ll \frac{r_{\text{int}}^2}{D_{3\text{d}}} \\ \frac{4\pi l_{\text{DNA}} D_{3\text{d}} t}{\ln(D_{3\text{d}} t / r_{\text{int}}^2)} & , t \gg \frac{r_{\text{int}}^2}{D_{3\text{d}}} \end{cases} \quad (14)$$

The first regime corresponds to reaction limited binding and is very brief for large $k_{\text{on}}^{\text{ns}}$. We discard this regime and ignore the slowly varying logarithm in the last regime, to approximate $P_{\text{surv}}^{\text{foreign}}$ by [21]

$$P_{\text{app}}(t) = \exp \left[-4l_{\text{DNA}} r_{\text{int}} \sqrt{\pi D_{3\text{d}} t} - k_{\text{cap}} t \right]. \quad (15)$$

The value of k_{cap} is fixed self-consistently by the condition that the total amount of proteins returning to the original segment is the same as for $P_{\text{surv}}^{\text{foreign}}$, i.e.

$$\int_0^\infty dt W_{\text{bulk}}^{\text{cyl}}(q, t) [P_{\text{surv}}^{\text{foreign}}(t) - P_{\text{app}}(t)] = 0. \quad (16)$$

We solved this equation numerically. But when a value for k_{cap} has been obtained then the choice of P_{app} in Eq. (15) is very convenient. In fact it can be written

$$P_{\text{app}}(t) = \int_0^\infty ds e^{-(s+k_{\text{cap}})t} \frac{a}{2\sqrt{\pi s^3}} e^{-a^2/(4s)}, \quad (17)$$

where $a = 4\sqrt{\pi} r_{\text{int}} l_{\text{DNA}} \sqrt{D_{3\text{d}}}$. When inserted into Eq. (13) in Laplace space we need to evaluate numerically only one integral to obtain $W_{\text{bulk}}(q, u)$, namely

$$W_{\text{bulk}}(q, u) = \int_0^\infty ds \frac{a e^{-a^2/(4s)}}{2\sqrt{\pi s^3}} W_{\text{bulk}}^{\text{cyl}}(q, u + s + k_{\text{cap}}). \quad (18)$$

The term with $\delta_{q,0}$ was already discarded here. Eq. (18) generalizes the straight rod result (9). In the limit $D_{3\text{d}} \gg k_{\text{on}}^{\text{ns}}$ it is consistent with the approximation $1 - \lambda_{\text{bulk}}(q) \approx 1$ leading to the expected result (10) for that case.

In above analysis we assumed that the density of foreign segments is uniform in space. Although this is not true for a relaxed coil (the density decreases on a scale of the persistence length of the DNA) we argue that this is not important for evaluating the search rate: in fact, if the protein diffuses far away from the original segment it will most likely perform a long relocation measured along the DNA contour. Again it does not matter for the search rate how we bookkeep these long relocations. To estimate the density l_{DNA} for a random relaxed coil we employ the Worm Like Chain model. A good approximation for the probability density for a point a contour distance s away from another point to loop back is [22]

$$j_M = \left(\frac{3}{4\pi|s|\ell_p} \right)^{3/2} \exp \left(-\frac{8\ell_p^2}{|s|^2} \right). \quad (19)$$

For a target in the middle of a chain of length L , we find the DNA density $l_{\text{DNA}} = \int_{-L/2}^{L/2} ds j_M$ around the target.

The theory presented here was shown in Ref. [16] to quantitatively describe recent experimental data obtained from an optical tweezers setup by which the search rate k_{on} at various degrees of DNA coiling could be measured. The relative increase R of k_{on} in the maximally relaxed DNA configuration compared to k_{on} in the stretched state, for DNA length $L = 6538$ base pairs, was measured to be around 1.1 to 1.3 in buffers with salt concentrations of 5 mM MgCl_2 and 0 to 25 mM for NaCl. This relative increase R compared well with the theory presented here using estimates of l_{DNA} based on Eq. (19) giving $1/\sqrt{l_{\text{DNA}}} \sim 500$ bp, and a value $l_{\text{sl}}^{\text{eff}} \sim 100$ bp based on a quantitative analysis [23] of experiments [24] suggesting effective sliding lengths of this size. Numerical analysis of the theory presented here showed that R depends significantly only on l_{DNA} and $l_{\text{sl}}^{\text{eff}}$ for realistic ranges of the remaining parameters. At physiological salt (100 mM NaCl) an increase of $R \approx 1.7$ was found experimentally [16]. This observation agrees well with the increased DNA density expected due to attraction between segments induced by the 5 mM divalent Mg^{2+} ions at this NaCl concentration, see Ref. [16] for details. We are not aware of a theory to calculate l_{DNA} under such attraction. However, the value $1/\sqrt{l_{\text{DNA}}} \sim l_{\text{sl}}^{\text{eff}}$ needed to obtain $R \approx 1.7$ at this salt conditions appears reasonable.

We presented a model for facilitated diffusion that is useful to calculate the search rate of DNA binding proteins for their specific binding site on a fluctuating DNA coil with variable DNA density. This model represents a convenient way to rephrase the Berg-von Hippel theory with the additional advantage that it interpolates between stretched and coiled DNA configurations. In the development of this theory a number of assumptions were made, in particular that of sufficiently fast local relaxation of the DNA configuration. This assumption was found to be satisfied for the protein EcoRV used in [16], but it will not hold universally. It would therefore be

beneficial to generalize the theory to slowly fluctuating or static DNA configurations.

We gratefully acknowledge discussions with Tobias Ambjörnsson, Alexander Grosberg, Mehran Kardar, Leonid Mirny, and Roland Winkler. This research was partially funded by the Villum Kann Rasmussen Foundation, Deutsche Forschungsgemeinschaft, and by a Fundamenteel Onderzoek der Materie Projectruimte Grant, a Netherlands Organization for Scientific Research-Vernieuwings Impuls grant.

-
- [1] B. D. Hughes *Random Walks and Random Environments*, Vol. 1 (Oxford University Press, Oxford, 1995).
 - [2] M. A. Lomholt, T. Koren, R. Metzler, and J. Klafter, *Proc. Natl. Acad. Sci. USA* **105**, 11055 (2008).
 - [3] T. Koren, M. A. Lomholt, A. V. Chechkin, J. Klafter, and R. Metzler, *Phys. Rev. Lett.* **99**, 160602 (2007).
 - [4] M. von Smoluchowski, *Physikal. Zeitschr.* **17**, 557 (1916).
 - [5] O. Bénichou, C. Loverdo, M. Moreau, and R. Voituriez, *Phys. Chem. Chem. Phys.* **10**, 7059 (2008).
 - [6] P.H. von Hippel and O.G. Berg, *J. Biol. Chem.* **264**, 675 (1989).
 - [7] I. M. Sokolov, R. Metzler, K. Pant, and M. C. Williams, *Biophys. J.* **89**, 895 (2005); J. Elf, G. W. Li, and X. S. Xie, *Science* **316**, 1191 (2007); Y. M. Wang, R. H. Austin, and E. C. Cox, *Phys. Rev. Lett.* **97**, 048302 (2006).
 - [8] A. Bakk and R. Metzler, *FEBS Lett.* **563**, 66 (2004); *J. Theor. Biol.* **231**, 525 (2004), and Refs. therein.
 - [9] O. G. Berg and M. Ehrenberg, *Biophys. Chem.* **15**, 41 (1982).
 - [10] Even in bacteria the DNA length measures several mm while DNA's persistence length is of the order of 50 nm.
 - [11] T. Hu, A. Y. Grosberg, and B. I. Shklovskii, *Biophys. J.* **90**, 2731 (2006).
 - [12] S. E. Halford and J. F. Marko, *Nucl. Acids Res.* **32**, 3040 (2004).
 - [13] M. Slutsky and L. A. Mirny, *Biophys. J.* **87**, 4021 (2004).
 - [14] M. Coppey, O. Bénichou, R. Voituriez, and M. Moreau, *Biophys. J.* **87**, 1640 (2004).
 - [15] M. A. Lomholt, T. Ambjörnsson, and R. Metzler, *Phys. Rev. Lett.* **95**, 260603 (2005).
 - [16] B. van den Broek, M. A. Lomholt, S.-M. J. Kalisch, R. Metzler, and G. J. L. Wuite, *Proc. Natl. Acad. Sci. USA* **105**, 15738 (2008).
 - [17] In practice the DNA should be longer than the effective sliding length $l_{\text{sl}}^{\text{eff}}$ introduced before Eq. (11).
 - [18] We deviate here from conventions in [16] by only letting n_{bulk} count 'active enzymes'.
 - [19] Eq. (11) generalizes Smoluchowski's result $k_{\text{on}} = 4\pi D_{3d}b$ for pure 3D search for a target of size b . The effective target size in Eq. (11) is often called antenna length.
 - [20] Note the distinction to intersegmental transfers where the protein intermittently is bound to two DNA segments [6].
 - [21] $P_{\text{surv}}^{\text{foreign}}$ and P_{app} agree very well numerically.
 - [22] L. Ringrose *et al.*, *EMBO J.* **18**, 6630 (1999).
 - [23] H.-X. Zhou, *Biophys. J.* **88**, 1608 (2005).
 - [24] A. Jeltsch and A. Pingoud, *Biochemistry* **37**, 2160 (1998); N. P. Stanford *et al.*, *EMBO J.* **19**, 6546 (2000).